

DAFTAR PUSTAKA

- Alfarisi, S. (2018). Pengaruh latihan PNF (Proprioceptive Neuromuscular Facilitation) dan thai massage terhadap tingkat fleksibilitas togok atlet UKM karate inkai UNY. In *Universitas Negeri Yogyakarta*.
- Ali, M. H., Gaikwad, S., & Mathew, J. T. (2015). A Comparative Study between Hold Relax Technique and Static Stretching to Improve Gait Parameter of Hemiplegic Stroke Patients. *International Journal of Neurorehabilitation*, 2(4), 1–4. <https://doi.org/10.4172/2376-0281.1000180>
- Carvalho, R., Dias, N., & Cerqueira, J. J. (2019). Brain-machine interface of upper limb recovery in stroke patients rehabilitation: A systematic review. *Physiotherapy Research International*, 1–16. <https://doi.org/10.1002/pri.1764>
- Chitra, J., & Joshi, D. (2017a). Effect of scapular hold-relax technique on shoulder pain in hemiplegic subjects: A randomized controlled trial. *Physiotherapy - The Journal of Indian Association of Physiotherapists*, 11(2), 49–52. https://doi.org/10.4103/pjiap.pjiap_11_17
- Chitra, J., & Joshi, D. D. (2017b). The effect of proprioceptive neuromuscular facilitation techniques on trunk control in hemiplegic subjects: A pre post design. *Physiotherapy - The Journal of Indian Association of Physiotherapists*, 11(2), 40–44. https://doi.org/10.4103/pjiap.pjiap_12_17
- Deb, P., Sharma, S., & Hassan, K. M. (2010). Pathophysiologic mechanisms of acute ischemic stroke: An overview with emphasis on therapeutic significance beyond thrombolysis. *Pathophysiology*, 17, 197–218. <https://doi.org/10.1016/j.pathophys.2009.12.001>
- Deepak Joshi, J. C. (2017). Effect of scapular proprioceptive neuromuscular facilitation on shoulder pain, range of motion, and upper extremity function in hemiplegic patients: A randomized controlled trial. *Indian Journal of Health Sciences and Biomedical Research KLEU*, 10(3), 276–282. <https://doi.org/10.4103/kleuhj.kleuhj>
- Dewi, N. V. K. (2016). *Stroke infark*.
- Hatem, S. M., Saussez, G., della Faille, M., Prist, V., Zhang, X., Dispa, D., & Bleyenheuft, Y. (2016). Rehabilitation of motor function after stroke: A multiple systematic review focused on techniques to stimulate upper extremity recovery. *Frontiers in Human Neuroscience*, 10, 1–22. <https://doi.org/10.3389/fnhum.2016.00442>
- Junior, V. A. dos S., Santos, M. de S., Ribeiro, N. M. da S., & Maldonado, I. L. (2019). Combining Proprioceptive Neuromuscular Facilitation and Virtual Reality for Improving Sensorimotor Function in Stroke Survivors: A Randomized Clinical Trial. *Journal of Central Nervous System Disease*, 11, 1–7. <https://doi.org/10.1177/1179573519863826>
- Kumar, B., Kumari, V. S., & Madhavi, K. (2013). The effect of Strength Training on Normalizing the Tone and Strength of Spastic Elbow Flexors in Subjects with Stroke. *Indian Journal of Physiotherapy and Occupational Therapy - An International Journal*, 7(3), 234. <https://doi.org/10.5958/j.0973->

5674.7.3.098

- Mumtazah, N., & Abdullah, F. (2020). Hold Relax dan Passive Stretching Efektif Dalam Meningkatkan Kemampuan Fungsional Pada Pasien Post-Gips Fracture Tibial Plateau Dextra. *Jurnal Ilmiah Fisioterapi*, 3(2), 16–23. <https://doi.org/10.36341/jif.v3i2.1397>
- Musik, T. T. (2011). *Anatomi Otak dan Fungsinya*.
- Nurhayati, U. A., Mahadewa, T. G. B., Lesmana, S. I., Krisna, I. M., Satriyasa, B. K., & Wahyudin. (2019). Proprioceptive neuromuscular fascilitation stretching lebih efektif dalam meningkatkan aktivitas fungsional daripada static stretching pada penderita osteoarthritis lutut. *Jurnal Fisioterapi Dan Rehabilitasi*, 3(2), 90–99.
- Ostwal, S., & Nagarwala, R. (2018). Effects of Neural Tissue Mobilisation and Pnf Technique of Hold-Relax on Lower Extremity Flexibility in Post Stroke Patients. *International Journal of Physiotherapy and Research*, 6(2), 2676–2681. <https://doi.org/10.16965/ijpr.2018.112>
- Ovbiagele, B., & Turan, T. N. (2016). *Ischemic Stroke Therapeutics A Comprehensive Guide*.
- Purwaningtyas, D. (2016). Pengaruh Pemberian Hold Relax terhadap Spastisitas pada Pasien Pasca Stroke. In *Universitas Hasanuddin Makassar*.
- Purwanto, H. (2016). *Keperawatan Medikal Bedah II*.
- Suwal, S., & N, P. K. B. (2020). A Study on the Effect of Lower Extremity Proprioceptive Neuromuscular Facilitation Patterns on Stair Ambulation in Stroke Patients. *Journal of Physiotherapy & Physical Rehabilitation*, 5(1), 1–4. <https://doi.org/10.37421/jppr.2020.05.181>
- Trompetto, C., Marinelli, L., Mori, L., & Pelosin, E. (2014). Pathophysiology of spasticity: Implications for neurorehabilitation. *BioMed Research International*, 1–8. <https://doi.org/10.1155/2014/354906>
- Ujwal Lakshman Yeole, Arora, S. P., Gharote, G. M., Panse, R. B., Pawar, P. A., & Kulkarni, S. A. (2017). Effectiveness of Proprioceptive Neuromuscular Facilitation on Spasticity in Hemiplegia: Randomised Controlled Trial. *Journal of Medical Science And Clinical Research*, 05(01), 15567–15572. <https://doi.org/10.18535/jmscr/v5i1.61>
- Wang, J. S., Lee, S. Bin, & Moon, S. H. (2016). The immediate effect of PNF pattern on muscle tone and muscle stiffness in chronic stroke patient. *Journal of Physical Therapy Science*, 28(3), 967–970. <https://doi.org/10.1589/jpts.28.967>
- Ward, A. B. (2012). A literature review of the pathophysiology and onset of post-stroke spasticity. *European Journal of Neurology*, 19, 21–27. <https://doi.org/10.1111/j.1468-1331.2011.03448.x>
- Widiarti, A. W., & Ratmawati, Y. (2018). Pengaruh pemberian “contrash bathing” terhadap penurunan spastisitas otot lengan dan tungkai pada anak dengan cerebral palsy. *Jurnal Terpadu Ilmu Kesehatan*, 7(1), 01–100.
- Wiley, J., Sons, & Ltd. (2013). *Organised inpatient (stroke unit) care for stroke (Review)* (Issue 9).
- Yeole, U. L., Arora, S. P., Gharote, G. M., Panse, R. B., Pawar, P. A., & Kulkarni, S. A. (2017). Effectiveness of Proprioceptive Neuromuscular Facilitation on

Spasticity in Hemiplegia: Randomised Controlled Trial. *Journal of Medical Science And Clinical Research*, 05(01), 15567–15572. <https://doi.org/10.18535/jmscr/v5i1.61>

Zhou, Z., Sun, Y., Wang, N., Gao, F., Wei, K., & Wang, Q. (2016). Robot-assisted rehabilitation of ankle plantar flexors spasticity: A 3month study with proprioceptive neuromuscular facilitation. *Frontiers in Neurorobotics*, 10, 1–14. <https://doi.org/10.3389/fnbot.2016.00016>