

DAFTAR PUSTAKA

- Alghadir, A. H., Anwer, S., Iqbal, A., & Iqbal, Z. A. (2018). *Test – retest reliability , validity , and minimum detectable change of visual analog , numerical rating , and verbal rating scales for measurement of osteoarthritic knee pain.* 851–856.
- Anwer *et al.*, (2014). *Effect of Isometric Quadriceps Exercise on Muscle Strength , Pain , and Function in Patients with Knee Osteoarthritis : A Randomized Controlled Study.* december, 745–748.
- Ariff. (2018). *Painful Pes Anserine Bursitis Following Total Knee Replacement Surgery.* 18(February), 97–99.
<https://doi.org/10.18295/squmj.2018.18.01.016>
- Cheon *et al.*, (2020). *Acute Effects of Open Kinetic Chain Exercise Versus Those of Closed Kinetic Chain Exercise on Quadriceps Muscle Thickness in Healthy Adults.* *International Journal of Environmental Research and Public Health*, 1–11.
<https://doi.org/10.3390/ijerph17134669>
- Curtis *et al.*, (2019). *Pes Anserinus : Anatomy.* *American Journal of Roentgenology*, 213(November), 1–10.
<https://doi.org/10.2214/AJR.19.21315>
- Enohumah. (2008). *Pain in Osteoarthritis.* *Afrika Journal of Biomedical Research*, 11(May), 119–128.
- Helfenstein, M., & Kuromoto, J. (2010). *Anserine syndrome.* 50(3), 320–327.
https://www.scielo.br/pdf/rbr/v50n3/en_v50n3a11.pdf
- Kim *et al.*, (2017). *Effects of open and closed kinetic-chain exercises on the muscle strength and muscle activity of the ankle joint in young healthy women.* *The journal of Physical Therapy Science*, VI, 1903–1906.
<https://doi.org/10.1589/jpts.29.1903>
- Kwon *et al.*, (2013). *The effect of open and closed kinetic chain exercises on dynamic balance ability of normal healthy adults.* *Journal of Physical Therapy Science*, 25(6), 671–674. <https://doi.org/10.1589/jpts.25.671>
- Lee *et al.*, (2014). *Pes anserinus and anserine bursa : anatomical study.* *Anatomy & Cell Biology*, 127–131. <https://doi.org/10.5115/acb.2014.47.2.127>
- Mesfar, W., & Shirazi-adl, A. (2008). *Knee joint biomechanics in open-kinetic-chain flexion exercises.* *Clinical Biomechanics*, 23(4), 477–482.
<https://doi.org/10.1016/j.clinbiomech.2007.11.016>

- Moghadam *et al.*, (2016). Comparing the Effect of Open and Closed Kinetic Chain Exercises in Patients Suffering From Patellofemoral Pain Syndrome. *International Journal of Basic Science in Medicine*, 1(2), 53–57. <https://doi.org/10.15171/ijbsm.2016.12>
- Mohseni. (2020). *pes anserine bursitis*. StatPearls.<https://www.ncbi.nlm.nih.gov/books/NBK532941/>
- Nejati, P., Farzinmehr, A., & Moradi-lakeh, M. (2015). The effect of exercise therapy on knee osteoarthritis : a random- ized clinical trial. *Medical Journal of the Islamic Republic of Iran*, 29.186, 1–9.
- Prio, Asminarsih Zainal., Misbah, Sitti Rachmi dan Wijayati, Fitri. (2017). Pengaruh Latihan Gerak Aktif Kaki Dengan Teknik Open Kinetik Chain Exercise Terhadap Penurunan Intensitas Nyeri Pada Lansia Dengan Nyeri Sendi Osteoarthritis Dan Rheumatoid. *Health Information : Jurnal Penelitian*. DOI: <https://doi.org/10.36990/hijp.v9i1.76> Juni (2017).
- Saggini *et al.*, (2015). Pes Anserine Bursitis in Symptomatic Osteoarthritis Patients : *Journal of Alternative and Complementary Medicine*, 21(8), 480–484. <https://doi.org/10.1089/acm.2015.0007>
- Sarifakioglu *et al.*, (2016). Comparison of the efficacy of physical therapy and corticosteroid injection in the treatment of pes anserine tendino-bursitis. *The Journal of Physical Therapy Science*, 28, 1993–1997. <https://doi.org/10.1589/jpts.28.1993>
- Waldman. (2008). *Pes Anserine Bursitis*. 468–471. <https://doi.org/10.1016/B978-0-323-54731-4.00118-3>
- Yarlagadda, S. D. (2013). A Study to compare the effect of Open versus Closed kinetic chain exercises in Patello-femoral arthritis . *IOSR Journal of Sports and Physical Education*, 1(1), 34–41.